

MicroRNA-146a controls local bone destruction by regulating fibroblast induced osteoclastogenesis in inflammatory arthritis

Victoria Saferding¹, Antonia Puchner¹, Eliana Goncalves-Alves¹, Melanie Hofmann¹, Julia Brunner², Emine Sahin², Silvia Hayer¹, Phillipe Georgel³, Marije M. Koenders⁴, Gernot Schabbauer², Josef S. Smolen¹, Günter Steiner¹, Kurt Redlich¹ and Stephan Blüml¹

¹ Division of Rheumatology, Internal Medicine III, Medical University of Vienna, Austria.

² Institute for Physiology, Center for Physiology and Pharmacology, Medical University Vienna, A-1090 Vienna, Austria.

³ INSERM UMR_S 1109, Fédération de Médecine Translationnelle (FM TS), Université de Strasbourg, Centre de Recherche en Immunologie et Hématologie, 1, Place de l'Hôpital 67085 Strasbourg Cedex France

⁴ Radboud University Nijmegen Medical Center, Nijmegen, The Netherlands

Background:

MicroRNA (MiR-) 146a plays an important role in the regulation of the innate immune response and has also been shown to suppress cancer development in myeloid cells. Although in late stages of arthritis elevated expression of miR-146a in synovial tissue of rheumatoid arthritis patients was detected, the level of this miRNA was found to be down regulated in early disease, but its role in the development of inflammatory arthritis is still elusive.

Materials and Methods:

To induce arthritis we used the chronic inflammatory hTNFtg disease model, therefore we crossed miR-146a deficient into hTNFtg mice. Disease severity was assessed clinically and histologically. Blood of arthritis animals was analysed by flow cytometry. Serum cytokine levels were measured by Elisa. Synovial fibroblasts were isolated from metatarsal bones. Proliferation of fibroblasts was analysed histologically and by ³[H]thymidine incorporation. RNA expression levels were measured by qPCR

Results:

When we crossed miR-146a^{-/-} into hTNFtg mice, histological examination revealed a significant increase in synovial inflammation and even more striking a more than twofold increase in local bone destruction, due to increased generation of osteoclasts in the tarsal joints in miR-146a^{-/-}/hTNFtg mice compared to hTNFtg mice. Interestingly, systemic bone loss was comparable in hTNFtg compared to miR-146a^{-/-}/hTNFtg mice, suggesting an important local role of miR-146a. Indeed, we detected increased levels of IL-1β, TRAF6, a major target of miR-146a and RANKL, in addition the expression level of OPG was decreased locally in the paws of miR-146a^{-/-}/hTNFtg compared to hTNFtg mice. By performing bone marrow transplants we could indeed show a pivotal role for miR-146a in mesenchymal cells in controlling local osteoclast generation and bone destruction. Analysis of important mesenchymal cells in arthritis, the synovial fibroblasts exhibited enhanced proliferation if miR-146a is missing, *in vitro* and *in vivo*. Moreover stimulation of these cells with IL-1β, a prominent cytokine in arthritis which was also shown to be negatively regulated by miR-146a, led to increased expression of RANKL and TRAF6 in miR-146a deficient synovial fibroblasts.

Conclusion:

These data demonstrate an important mitigating role of the miR-146a in inflammatory arthritis, most importantly in local bone destruction, by controlling mesenchymal expression of osteoclastogenic factors. This shows an important anti-inflammatory role of miR-146a, which might possibly be exploited for therapeutic purposes.